

**READ ENTIRE LABEL BEFORE EACH USE
FOR PET USE - USE ONLY ON CATS OR DOGS**

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Humans: Harmful if swallowed or if inhaled. Remove contaminated clothing and wash clothing before reuse. Avoid breathing dust. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. **Animals:** Harmful if swallowed, inhaled. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals or animals on medication. Sensitivities may occur after using **ANY** pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact SafetyCall International (866) 899-8229 for emergency medication information.

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

USE ON LIVESTOCK

BEEF AND DAIRY CATTLE AND HORSES
Can be used in dust bags, shaker cans and mechanical dust applicator.

Horn Flies, Lice, Face Flies: Place contents of this package in any commercially available dust bag, suspend bag in area frequented by cattle or in gateway or lanes through which the animals must pass daily for water, feed or minerals. Bags may also be placed in feeding sheds or in front of mineral feeders.

For dairy cows, bags may be suspended in the neck through which the cows leave the milking barn. The bags should hang 18 to 24 inches below the back line of the cattle. For reduction of Face Flies, bags must be located so animals will be forced to use them daily and hung at a height so that the face is dusted.

Horn Flies, Lice Direct Application: Apply 2 oz. of dust per animal to shaker can over the head, neck, shoulders and back and behind. Repeat as necessary, but not more than once every three days.

SWINE:
Lice on Swine Direct Application: Apply only 1 ounce per head as a uniform coat to the head, shoulder and back by use of a shaker can or suitable mechanical dust applicator. Repeat as necessary, but not more often than once every 10 days. In severe infestation, both animals and the bedding may be treated. Do not ship animals for slaughter within 5 days of treatment.

POLYTRIX: To control Northern Fowl Mites, apply at a rate of 1 lb. per 100 birds. Ensure thorough treatment of vent area.

USE ON CATS AND DOGS

To control Fleas, Ticks and Lice on Cats and Dogs:
Use product only outside or in a well-ventilated area. Wearing household latex or rubber gloves dust animals thoroughly, working the dust into hair down to the skin. Pay close attention to legs and feet. Use 1/2 oz. on animals 20 pounds and under and 1 oz. on pets over 20 pounds. Avoid getting dust in pet's eyes, nose, mouth and genital areas. Repeat every three weeks as needed. Do not use on cats or dogs under twelve (12) weeks of age. Dust sleeping area including bedding and cracks and crevices.

STORAGE AND DISPOSAL
PESTICIDE STORAGE: Store in original container. Keep area inaccessible to children and pets.
DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Do not reuse unless required for recycling. If recycling is not available then dispose of container in a suitable landfill or if allowed by State and local authorities, by burning. If burned, stay out of smoke.
If partly filled: Call your local solid waste agency for disposal instructions. Never pour unused product down any indoor or outdoor drain or on the ground.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic organisms. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwater. This pesticide is toxic to honey bees and other beneficial pollinators exposed to an application. Do not apply when bees are actively visiting blooming plants (vegetables, flowers, Frustrationment trees). Do not allow this dust to come in direct contact with bees at any time.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOME GARDENS: FOR USE ON HOME GARDENS

NOT FOR USE ON PLANTS BEING GROWN FOR SALE OR OTHER COMMERCIAL USE.

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous dusting rotary duster or other appropriate dusting as available. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. (Many insects hide on the underside of the leaf.)

Apply when air is calm to avoid drift and contact with eyes and skin. Controls a wide range of insect pests. Use at first signs of insect infestation, preferably before damage from feeding occurs. **Note the pre-harvest time between use and harvest interval with each crop listed.**

VEGETABLES: DIRECTIONS FOR APPLICATION

CROP	TARGET PEST	REMARKS
Asparagus	Cutworm Complex, Asparagus Beetle	Dust lightly. Apply up to 4 times per season. May be applied until 1 day before harvest.
Broccoli, Brussels Sprouts, Cauliflower	Cabbage Looper, Imported Cabbageworm, Diamondback Moth, Cabbage Aphid	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.
Cabbage	Cabbage Looper, Imported Cabbageworm, Diamondback Moth, Cabbage Aphid	Dust lightly. Apply up to 5 times per season. May be applied until 1 day before harvest.
Cucumber, Muskmelons, Pumpkins, Squash, Watermelon	Leafminer, Squash Bug, Cabbage Looper, Cucumber Beetle (adult), Cutworm, Melon Worm, Pickle Worm, Plant Bug (including signs), Red Sinker Bug, Broodworm, Squash Vine Borer	Dust lightly. Apply up to 8 times per season. May be applied until harvest.
Celery, Lettuce (head and leaf), Parsley, Rhubarb, Spinach	Vegetable Leafminer, Cabbage Looper, Cutworm Complex, Fall Armyworm, Beet Armyworm, Corn Earworm, Leafhopper, European Corn Borer	Dust lightly. Apply up to 5 times per season. May be applied until 1 day before harvest.
Eggplant	Colorado Potato Beetle, Vegetable Leafminer, Cabbage Looper, Flea Beetle	Dust lightly. Apply up to 10 times per season. May be applied until 3 days before harvest.
Onions (Dry bulb and Garlic)	Onion Thrips, Armyworm, Cutworm, Leafminer, Onion Maggot (adult), Slink Bug	Dust lightly. Apply up to 6 times per season. May be applied until 1 day before harvest.
Peppers (Bell)	Pepper Weevil, Vegetable Leafminer, Cabbage Looper, Flea Beetle, European Corn Borer	Dust lightly. Apply up to 8 times per season. May be applied until 3 days before harvest.
Potatoes	Colorado Potato Beetle, Potato Leafhopper, Aster Leafhopper, Potato Flea Beetle, Potato Aphid, Potato Tuberworm, Potato Psyllid, Cabbage Looper	Dust lightly. Apply up to 8 times per season. May be applied until 14 days before harvest.
Sweet Corn	Corn Earworm, European Corn Borer, Fall Armyworm, Southern Armyworm, Corn Rootworm (adult), Cutworm, Aster Leafhopper, Flea Beetle	Dust lightly. Apply up to 6 times per season. May be applied until 1 day before harvest.
Tomatoes	Tomato Fruitworm, Cabbage Looper, Tomato Pinworm, Vegetable Leafminer, Beet Armyworm, Southern Armyworm, Colorado Potato Beetle, Granulate Cutworm, Hornworm	Dust lightly. Apply up to 6 times per season. May be applied until day of harvest.

ORNAMENTALS: DIRECTIONS FOR APPLICATION

CROP	TARGET PESTS	REMARKS
Asparagus Fern	Japanese Beetle, Asparagus Beetle, Lygus Bug	Dust lightly. Apply to the fern stage of asparagus plant after spear harvest when adult stage of the asparagus beetle, lygus bugs and the adult stage of the Japanese beetle are present.
Roses	Beet Armyworm, Cabbage Looper, Omivirous Leafhopper	Dust lightly.

TREE CROPS: DIRECTIONS FOR APPLICATION

CROP	TARGET PESTS	REMARKS
Apples	Apple Aphid, Redbanded Leafhopper, Obliquebanded Leafhopper, Plum Curculio, White Apple Leafhopper, Spotted Tentiform Leafminer, Tarnished Plant Bug	Dust lightly. Do not apply after petal fall.
Peaches	Peach Twig Borer, Oriental Fruit Moth, Green Fruitworm, Lesser Peachtree Borer, Plum Curculio, Tarnished Plant Bug	Dust lightly. Apply up to 8 times per season. May be applied until 14 days before harvest.
Walnuts	Navel Orangeworm, Codling Moth, Walnut Husk Fly	Dust lightly. Apply up to 8 times per season. May be applied until 1 day before harvest.

WARRANTY STATEMENT


Read the entire Directions for Use and Warranty before using this Product. By using this product, user or buyer accepts the following warranty. The directions for use of this product are believed to be adequate and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Unintended consequences may result because of unknown factors. All such risks shall be assumed by the user or buyer. Control Solutions Inc. is committed to providing high quality products. To the extent consistent with applicable law, Control Solutions Inc. makes no warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. To the extent consistent with applicable law, Control Solutions Inc., the manufacturer, or the Seller shall not be liable for indirect, special, incidental or consequential damages. To the extent consistent with applicable law, the exclusive remedy of the user or buyer shall not exceed the purchase price paid.

EPA Reg. No. 53883-33 EPA Est. No. 44616-MO-1

Control Solutions, Inc.
5903 Geneva Road
Pasadena, TX 77507-1041
Phone (281) 892-2500



NET CONTENTS: 4 POUNDS



Easy-to-Use Applicator Nozzle

A CONTROL SOLUTIONS INC. PRODUCT

See Additional Precautionary Statements on Back of Bag.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

ACTIVE INGREDIENT:
Permethrin (3-phenoxyphenyl) methyl (+/-) cis, trans-3-(2,2-dichloroethenyl)-2, 2-dimethylcyclopropanecarboxylate* 0.25%
OTHER INGREDIENTS: 99.75%
TOTAL: 100.00%
* cis/trans ratio: Min. 35% (+/-) cis Max. 65% (+/-) trans

For Outdoor Use on Home Gardens
• For use on poultry, dogs, cats, horses, swine, beef and dairy cattle
• Controls A Wide Range of Insect Pests On Garden Vegetables
• Kills Over 60 Insects on Vegetables, Ornamentals and Fruits
As Listed

VIPER INSECT DUST

CONTROLS:





12 x 4 LB.

VIPER INSECT DUST

Control Solutions, Inc.
Pasadena, TX 77507



Product Code MART4017